

EVALUATION OF BIOACTIVE CONSTITUENTS PRESENT IN *PSORALEA CORYLIFOLIA*

Tiwari, P.

Assistant Professor,
Department of Botany,
Shri Shivaji Science College, Congress Nagar, Nagpur-440012.

Abstract

Plants produce a diverse range of bioactive molecules making them a rich source of different type of medicines. The active complementary components give the plant as a whole a safety and efficiency much superior in the field of medicine and health. *Psoralea corylifolia* as one of the medicinal herbs is used in the treatment of many diseases and in many medicinal formulations. In present investigation, the phytochemical content of *P. corylifolia* is studied using leaf and stem as source material in two different organic solvents i.e. petroleum ether and methanol extracts.

Keywords: *P. corylifolia*, phytochemical screening, organic solvents

Introduction:

Nature is the biggest source of wonderful gifts to the human welfare. The importance of herbs in the management of human ailments cannot be overemphasized. The active components of herbal remedies have the advantage of being combined with many other substances that appear to be inactive(1). However, these complementary components give the plant as a whole a safety and efficiency much superior to that of its isolated and pure active components (2). *Psoralea corylifolia* as one of the medicinal herbs is used in the treatment of many diseases and in many medicinal formulations (3). *Psoralea corylifolia* L. [Indian breadroot] commonly known as Babachi is reported as a rare and endangered plant species which is available in tropical and subtropical regions of the world.(4) Numerous studies have identified compounds within herbal plants that are effective antibiotics(5). The screening of plant extracts and plant products for antimicrobial activity has shown that higher plants represent a potential source of novel antibiotic prototypes(6).

Psoralea is one of the main herbs in traditional Indian and Chinese herbal medicine for the treatment of skin disorders. It has been used in the treatment of eczema and hair loss. Roots of the plant are useful in dental caries, fruits are laxative, aphrodisiac, and are used for the treatment of leucoderma, leprosy and in inflammatory diseases of the skin and leaves are good for the treatment of diarrhoea. The plant has been used in Ayurvedic medicinal system as a cardiac tonic, vasodilator and pigmentor. (7). The 250,000-300,000 species of higher plants

were the main sources of drug for the world population. Much attention have been focused on phytochemicals as potential sources of functional substances (8) such as antioxidants(9), antiplague substances (10), antimutagenicities (11), enzyme inhibitors (12) and antimicrobial substances (13-15).

Aqueous and alcoholic extracts from *Psoralea corylifolia* leaves were screened and revealed the presence of saponins, tannins, flavonoids, glycosides, carbohydrates, tannins and phenolic compounds, gums and mucilages, fixed oils and fats. *P. corylifolia* leaves in respect to their antimicrobial activity and the broad spectrum of activity makes it a promising indigenous drug (16). Screening of babchi oil showed that the essential oil contain most of the phytochemicals including tannins, glycosides, saponins, flavonoids, steroids, terpenoids and flavonosides. (7). *Psoralea corylifolia* is a major contributor in the field of health science. A lot of research work is being carried on it due to aspects of its phytochemical contents.

Materials and Methods:

Plant materials:

Leaves and tender stem of *Psoralea corylifolia* were obtained from the plants grown in the college garden. Seeds were purchased from the local medicinal plant agency in Nagpur city. Leaves, tender stem and seeds were washed, air dried and then powdered in mixed grinder and stored in air tight bottles.

Preparation of solvent extracts:

Solvent extracts of *P. corylifolia* were prepared by using leaves and stem as a plant material. Two different solvent extracts used for phytochemical study were methanol and

petroleum ether prepared in cold and hot extraction method. Standard protocols were followed for studying phytochemical content of test material as given in Table no.1. Solvent extraction was prepared by taking 25grams of powder in 200ml of solvent in a conical flask. For best extraction, a soxhlet extractor was used for 48 hours. After this, extracts was concentrated through rotator evaporator which was then stored at 4°C (8).

Phytochemical screening of plant extracts:

A standard protocol is followed to detect the presence of various contents present in different parts (leaf and stem) of *P.corylifolia* as given in Table no.1.

Result and Discussion:

In present investigation, the phytochemical content of *P.corylifolia* is studied using leaf and stem as source material. Among fifteen different contents tested with their standard phytochemical methods given in Table no.1.

Aucubins and iriodoids, coumarins and fatty acids/lipids were detected in both the solvent extracts (methanol and petroleum ether). Table no.2 depicts the presence and absence of different bioactive compounds detected using their respective methods.

Babchi oil dissolved in methanol was evaluated for the presence of different phytochemicals to ascertain the presence of metabolites such as reducing sugars, alkaloids, anthraquinones,

glycosides, flavonoids, tannins, steroids, saponins, triterpenoids and phlobatanins. However, anthraquinone, phlobatanins and reducing sugars were not observed in babchi oil. Somasundaram *et al.*, 2010 reported the presence of saponins, tannins, flavonoids, glycosides, carbohydrates, tannins and phenolic compounds, gums and mucilages, fixed oils and fats in *Psoralea corylifolia* leaf extracts. Alcoholic extract is better than that of aqueous extract of *P. corylifolia* leaves in respect to their antimicrobial activity and the broad spectrum of activity makes it a promising indigenous drug.

Conclusion:

Plants produce a diverse range of bioactive molecules making them a rich source of different type of medicines. There is a great need to focus on the phytochemical study of plants to use them for more and more therapeutic purposes. In the present study, the approach is done to study the phytochemical evaluation of *P. corylifolia* in two different extracts.

Acknowledgment:

The author is thankful to Department of Botany, Science College, Congress Nagar, Nagpur, to use the lab and equipments to perform the research and also thankful to Minal Pathak, Research scholar to assist in the study sincerely.

References:

1. Parekh, J. and Chanda, S., 2007, *In vitro* antibacterial activity of the crude methanol extract of *Woodfordia fruticosa* Kurz. flower (Lythaceae). Braz. J. Microbiol, 38, 204-207.
2. Shariff, Z.U., 2001, Modern Herbal Therapy for Common Ailments. Nature Pharamcy Series Vol.1, Spectrum Books Ltd., Ibadan, Nigeria in Association with Safari Books (Export) Ltd. UK, 9-84.
3. Baikar, S. and Malpathak, N., 2011, Genetic characterization and biochemical analysis in the populations of *P. corylifolia* L. from Western Maharashtra. J.of Biodiversity and Eco.Sci, 1(4), 280-288.
4. Kaufman, P. Duke, P. Brielmann, J. Boik, H. and Hoit, J., 1997, Comparative survey of leguminous plants as a source of isoflavones genistein and daidzein implication for human nutrition and health. J. Alter comp Med, 3(1), 7-12.
5. Basile, A. Sorbo, S. Giordano, S. Ricciardi, L. Ferrara, S. Montesano, D. Castaldo Cobianchi, R. Vuotto, M.L. and Ferrara, L., 2000, Antibacterial and allelopathic activity of extract from *Castanea sativa* leaves. Fitoterapia, 71, 110-116.
6. Afolayan, A.J., 2003, Extracts from the shoots of *Arctotis artotoides* inhibit the growth of bacteria and fungi. Pharm. Biol, 41, 22-25.
7. Purkayastha, S. and Dahiya, P., 2012, Phytochemical screening and antibacterial potentiality of essential oil of *Psoralea corylifolia* L. *Inter.J. of Bio Sci., Bio Chem. And Bio Info*, 2(3), 188-191.

8. Chanda, S. Kaneria, M. and Nair, R., 2011, Antibacterial Activity of *Psoralea corylifolia* L. Seed and Aerial Parts with Various Extraction Methods. Research Journal of Microbiology, 6, 124-131.
9. Esaki, H. Watanabe, R. Onozaki, H. Kavakishi, S. and Osawa, T., 1999, Formation mechanism for potent antioxidative o-dihydroxyisoflavones in soyabeans fermented with *Aspergillus saitoi*. Biosci. Biotechnol. Biochem, 63, 851-858.
10. An, B. J. Bae, M.T. and Choi, C., 1998, Chemical structure and isolation of glucosyltransferase inhibitor from the leaves of Korean persimmon. Food Sci. Biotechnol, 7, 23-27.
11. Karakaya, S. and Kawas, A., 1999, Antimutagenic activity of some foods. J. Sci. FoodAgric, 79, 237-242.
12. Choi, S.W. Kim, H.J. Chang, E.T. and Sapers, G.M., 1997, Inhibition of tyrosinase activity by plant extracts. Food Biotechnol, 6, 44-49.
13. Yim, C.K. Moon, J.H. and Park, K.H., 1999, Isolation of 3,4 dihydroxybenzoic acid, which exhibits antimicrobial activity, from fruits of *Gardenia jasminoides*. Kor. J. Food Sci. Technol, 31, 1386-1392.
14. Rojas, R. Bustamante, B. Bauer, J. Fernandez, A. Alban, J. and Lock, O., 2003, Antimicrobial activity of selected Peruvian medicinal plants, J. Ethnopharmacol, 88, 199-204.
15. Duraipandiyar, V. Ayyanar, M. and Ignacimuthu, S., 2006, Antimicrobial activity of some ethnomedicinal plants used by Paliyar tribe from Tamil Nadu, India. BMC Complement, Altern. Med, 6, 35-35.
16. Somasundaram, T. Chandrasekar, S.B. Bhanumathy, M. and Vijay, A.R., 2012, Phytochemical investigation and antimicrobial studies on the leaf extracts of *Psoralea corylifolia* Linn, Phcog. J, 4 (13), 51-55.

Table no.1: Phyto-chemical screening test of *P.corylifolia*

Compound	Compound	Reagent	Colour observation
Alkaloids	P.ether, chloroform, acetone, Ethanol, water extract	5ml 15% HCL Wagners reagent Mayers reagent Dragendroff reagent	Brown flocculent ppt., yellowish white ppt., orange ppt.
Aucubins/ Iriodoids	Fresh plant material	Trim Hill reagent	Blue color, green and red color
Cardiac glycosides	Water extract	Kedde,s test Legal test	Blue/violet color Pink color
Coumarins	P.ether chloroform, acetone, Ethanol, water extract	Residue+ P. ether Filterpaper moistened with 10% w/v aq.NaOH	Filter paper observed UV light shows yellow green flourescent
Tannins	Alcoholic and water extract	0.5ml extract+1ml H2O+2-3drops dilute 10% w/v aq. ferric chloride solution	Blue or green black color
Anthocyanins/ Anthocyanidines	Alcoholic and water extract	pH3-4 pH8-9	Red color changes after change in pH
Anthracene Glycosides	Alcoholic and water extract	Ethereal solution+25% v/v aq.NH4OH	Red color
Carotenoids	P.ether	Conc.HCL+Phenol	Blue or green color
Cynogenic Glycosides	Fresh plant material	Few drops of chloroform filter paper moistened with Na picrate solution	Filter paper turns red
Steroids	P.ether chloroform, acetone, Ethanol, water extract	Salkowski reaction Burchard reaction	Red color chloroform layer and acidic lower layer gives yellow flourescence
Emodins	P.ether	Borntragers reaction+benzene +25% w/v aq.NH4OH	Red color
Fattyacids/ Lipids	P.ether	Filter paper	Translucent spot on filter paper
Flavonoids	P.ether chloroform, acetone, Ethanol, water extract	Shinodas reaction Residue of ethereal solution+ethanol+ Mg.powder+conc.HCL	Red color
Triterpenoids	P.ether chloroform, acetone, Ethanol, water extract	Libermann- Burchard reaction	Red or violet color
Anthraquinones	Fresh plant material	Extracted with .55% w/v KOH +1ml H2O2+1ml acetic acid +1ml benzene+equal volume of dilute NH4	Red color in ammonical layer

Table no.2: Presence (+) and absence (-) of active compounds in *P.corylifolia* in different solvents during phytochemical screening.

Compound	Petroleum ether (HOT EXTRACT)		Petroleum ether (COLD EXTRACT)		Methanol (HOT EXTRACT)		Methanol (COLD EXTRACT)	
	Leaf	Stem	Leaf	Stem	Leaf	Stem	Leaf	Stem
Alkaloids	-	-	-	-	+	-	+	-
Aucubins/ Iriodoids	+	+	+	+	+	+	+	+
Cardiac glycosides	-	-	-	-	-	-	-	-
Coumarins	+	+	+	+	+	+	+	+
Tannins	-	-	-	-	+	-	+	-
Anthocyanins/ Anthocyanidines	-	-	-	-	-	-	-	-
Anthracene Glycosides	-	-	-	-	-	-	-	-
Carotenoids	+	-	-	+	+	-	+	-
Cynogenic Glycosides	-	-	-	-	-	-	-	-
Steroids	+	-	+	-	-	-	-	-
Emodins	-	-	-	-	-	-	-	-
Fattyacids/ Lipids	+	+	+	+	+	+	+	+
Flavonoids	-	-	-	-	-	-	-	-
Triterpenoids	-	-	-	-	-	-	-	-
Anthraquinones	-	-	-	-	-	-	-	-