

# Mortality of Tasar Silk worm, *Antheraea mylitta* D. Caused by Insect Predators

**B. S. Rahile**

Science College Pauni, Distt. Bhandara (M.S.)

## Abstract:

Wild tropical tasar silk worm, *Antheraea mylitta* D. reared in forest adjoining area mainly in four districts of eastern Vidarbha; Gadchiroli, Chandrapur, Bhandara and Gondia. It is regarded as a symbol of tribal culture and has been associated with many tribal group such as; Bhill, Gond, Rajgond, Laman, Varli, Katakare, Pradhan and Dhiwar. In the present study the severity and incidences of insect predators of tasar silk worm, *Antheraea mylitta* D. were studied during year 2010-2013 at three different tasar growing locations of Vidarbha of Maharashtra during different tasar rearing crops. The mortality of tasar silk worm due to insect predators was found to vary during different seasons of the year at different locations. Most of the damage was caused by the hemipteran reduvid bug, *Sycanus collaris*; stink bug, *Cantheconafurcellata*; Common wasps, *Vespa orientalis* and praying mantis, *Hierodullabipapilla*.

**Keywords:** Tropical tasar silk worm, *Antheraea mylitta*, insect predators, reduvid bug, stink bug, common wasps, praying mantis

## Introduction:

Tropical tasar silk worm, *Antheraea mylitta* D. is a lepidopteran holo- metabolous multivoltine and polyphagous insect. In exothermic insects, the environmental factors such as temperature, humidity, rainfall etc. play important role in growth and development in every stage (Back, 1980). Due to outdoor nature of rearing of tasar silk worm, is exposed to various rigors of the changing environment during different rearing seasons of the year, which often leads to outbreak of diseases and pest attack in rearing affecting the productivity. The predators of tasar silk worm are natural enemies in abundance in the rearing field and cause crop loss up to 20-25% (Singh et al.1992).The present study was carried out during the year 2009 to 2012 in three identified villages viz. Nisti, Tadgaon and Armori of Bhandara, Gondia and Gadchiroli district of Maharashtra respectively to study the mortality of tasar silk worm, *Antheraea mylitta* D. caused by insect predators in different rearing cycles.

## Materials and Methods:

The study was carried out in the tasar rearing fields one in each of districts Bhandara, Gondia and Gadchiroli of Maharashtra during year 2010 to 2013. The seed crops reared during the months of July- August and September- October and commercial crops during the months of November –January were considered for the study. In each crop attack of different insect pests was recorded. All the meteorological parameters like temperature, relative humidity and rainfall corresponding to different crops were recorded to study the correlation with incidences of different pest attack.

## Results and Discussion:

Three year pooled data pertaining to crop and place wise incidences of insect predators in relation to temperature, relative humidity and rainfall were collected and presented in Table 1 and 2. Variations in temperature can affect the biology, later searching ability and rate of parasitism and predation (Mack *et al*, 1981; Flinn, 1991) and the functional response (Messenger, 1989). Variation in development time increased with decreasing temperature and larger than expected based on the general relationship for insects and mites as observed by Shaffer (1983). In the present study the predators were observed in the month of July and the population reduced in month of August. They reappeared again in September and persisted upto October. The cycle of these insect predators was well



synchronized with the cycle of tasar silkworm. The incidence was high during third crop might be due to prolongation of larval duration in winter. The result of present study is in accordance with previous observation made by Shaffer (1983) reported that in *A. calandreae* females, longevity was longer at lower temperature and shortest at higher temperature. Whereas Islam *et al.*, (2005) reported that the development period reduced with increased temperature in *Dinamusbasalis*.

In the present study, most of the damaged caused by the hemipteran sting bug followed by wasps, mantis and reduvidbug (Fig. 2) during all the three cycles. The mortality was high due to *Canthecona* (Fig. 1) as the single bug can kill about 130-220 tasar larvae by sucking haemolymph of young instars. Senet. *al.*, (1971) reported same trend in mortality of tasar silkworm by stink bug. Even though wasps (Fig. 3) and mantis (Fig. 4) were observed during all the three cycles the mortality was low since they attacked mostly young age larvae, and young age rearing was undertaken in the controlled conditions at young age rearing sites.

**Table.1-** Incidences of insect pests in tasar silkworm, *Antheraea mylitta* D. during different seasons at different rearing sites (in percent)

Location	Insect Pests	Mortality		
		June-July (First Crop)	Aug.-Sept. (Second Crop)	Nov. -Jan. (Third Crop)
Tadgaon (Gondia)	Wasp	4.12	3.12	2.19
	Praying Mantis	2.87	2.38	3.65
	Reduvid Bug	1.00	2.06	0.83
	Stink Bug	5.18	8.89	4.53
Nisti (Bhandara)	Wasp	4.13	3.09	2.71
	Praying Mantis	4.00	5.23	6.11
	Reduvid Bug	1.20	3.46	1.14
	Stink Bug	5.14	6.12	6.20
Armori (Gadchiroli)	Wasp	5.20	3.09	2.20
	Praying Mantis	5.37	5.51	3.28
	Reduvid Bug	1.85	3.00	0.91
	Stink Bug	7.43	8.36	4.60

**Table. 2-** Weather parameters in different locations during the occurrence and incidences of insect pests in tasar silkworm, *Antheraea mylitta* D.

Location	Weather Factor	June-July	Aug.-Sept.	Nov. -Jan.
Tadgaon (Gondia)	Max. Temp. ( $^{\circ}$ C )	40.5	36.1	33.1
	Mini.Temp. ( $^{\circ}$ C )	22.4	22.2	8.6
	Max. RH (%)	100	100	96
	Mini. RH (%)	31	46	28
	Rainfall (mm)	341	268	22
Nisti (Bhandara)	Max. Temp. ( $^{\circ}$ C )	42	36.4	33.1
	Mini.Temp. ( $^{\circ}$ C )	22	22	9.5
	Max. RH (%)	96	96	88
	Mini. RH (%)	32	43	29
	Rainfall (mm)	338	302	50
Armori (Gadchiroli)	Max. Temp. ( $^{\circ}$ C )	41	36.5	33.6
	Mini.Temp. ( $^{\circ}$ C )	22	22.2	10
	Max. RH (%)	98	96	90
	Mini. RH (%)	30	56	25
	Rainfall (mm)	286	226	17



**Figure. 1-** Stink bug, *Cantheconafurcellata* **Figure. 2-** Reduvid bug, *Sycanuscollarissucking* haemolymph from tasar silkworm larvae



**Figure. 3-** Colony of common wasps, *Vespa orientalis* **Figure. 4-** Praying mantis, *Hierodullabipapilla*

## References

- Beck, S. D. (1980)** Insect Photoperiodism, Academic press, New York, II Edition.
- Flinn, P.W. (1991).** Temperature dependence functional response of *Cephalonomiawaterstoni* (Hymenoptera: Bethylidae) attacking rusty grain beetle larvae. *Environmental Entomology*, 20: 872-876.
- Islam, N., Islam, W. and Mondal, K. A. M. S. H. (2005)** Effect of temperature on life history characteristics of *Dinarmusbasalis* (Rond.) (Hymenoptera:Pteromalidae), a parasitoid of *Callosobruchus maculates* (F.) *Entomon* 30 (1):47-55.
- Mack, T. P., Bajusz, B. A., Nolan, E. S. and Smilowitz, S. (1981).** Development of a temperature mediated functional response equation. *Environmental entomology*, 10: 573-579.
- Messenger P. S. (1989).** Bioclimatic study of the aphid parasite *Praonexoletum*. Effects of temperature on the functional response of females to varying host densities. *Canadian Entomology* 100: 728-740.
- Sen, S.K, Jolly, M.S. and Jammy, T.R. (1971)** Biology and life cycle of *C. furcellata* Wolff (Hem: Pentatomidae) predator of tasar silkworm *A. mylitta* D. *Indian J. Seric.*, 10: 53-56.
- Shaffer, P.L. (1983)** Prediction of variation in development period of insects and mites reared at constant temperatures. *Environmental entomology*, 12: 1012-1019
- Singh R.N., Bajpayee C.M., Jayaswal J. and Thangavelun (1992)** Perspective of biological control in tasar culture, *Indian silk*, 31 (7): 48-50